

Pneumonia

| Key Indicators | Holy Cross Hospital (1) | National Average (2) | MD State Average (2) | Top 10% of Hospitals Nationwide (2) |
|--|-------------------------|----------------------|----------------------|-------------------------------------|
| Percent of Pneumonia Patients Assessed and Given Pneumococcal Vaccination | 98% | 87% | 88% | 100% |
| Percent of Pneumonia Patients Whose Initial Emergency Room Blood Culture Was Performed Prior To The Administration Of The First Hospital Dose Of Antibiotics | 97% | 92% | 91% | 100% |
| Percent of Pneumonia Patients Given Smoking Cessation Advice/Counseling | 100% | 91% | 95% | 100% |
| Percent of Pneumonia Patients Given Initial Antibiotic(s) within 6 Hours After Arrival | 96% | 94% | 94% | 100% |
| Percent of Pneumonia Patients Given the Most Appropriate Initial Antibiotic(s) | 94% | 89% | 91% | 99% |
| Percent of Pneumonia Patients Assessed and Given Influenza Vaccination | 98% | 85% | 83% | 100% |

*The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.

(1) Trinity Health data time period: October 2009 to March 2010.

(2) National average and Top 10% of hospitals nationwide were obtained from the Hospital Compare website on July 21, 2010. The latest reporting period is from October 2008 through September 2009 for most indicators.